arXiv:1302.5737v1 [hep-ph] 22 Feb 2013

Coulomb gauge model for hidden charm tetraquarks

W. Xie[a], L. Q. Mo[a], Ping Wang[ab], Stephen R. Cotanch[c]

[a] *Institute of High Energy Physics, CAS, P. O. Box 918(4), Beijing 100049, China*

[b] *Theoretical Physics Center for Science Facilities, CAS, Beijing 100049, China*

[c] *Department of Physics, North Carolina State University, Raleigh, NC 27695-8202, USA*

PACS: 12.39.Mk; 12.39.Pn; 12.39.Ki; 12.40.Yx; 03.70.+k; 11.10.-z; 11.10.Gh

Keywords: Exotic mesons, hidden charm, tetraquarks, QCD Coulomb gauge, effective Hamiltonian

---

Abstract

The spectrum of tetraquark states with hidden charm is studied within an effective QCD Coulomb gauge Hamiltonian approach. Of the four independent color schemes, two are investigated, the $(q\bar{c})_1(c\bar{q})_1$ singlet-singlet (molecule) and the $(qc)_3(\bar{q}\bar{c})_3$ triplet-triplet (diquark), for selected J^{PC} states using a variational method. The predicted masses of triplet-triplet tetraquarks are roughly a GeV heavier than the singlet-singlet states. There is also an interesting flavor dependence with $(q\bar{q})_1(c\bar{c}_1)$ states about half a GeV lighter than $(q\bar{c})_1(\bar{q}c)_1$. The lightest 1^{++} and 1^{--} predictions are in agreement with the observed $X(3872)$ and $Y(4008)$ masses, respectively, suggesting they both have a $\omega J/\psi$ rather than $D^*\bar{D}^*$ molecular type structure. This is consistent with observed X, Y decays to $\pi^{\pm}J/\psi$ and not $D^*\bar{D}^*$.

---

1 Introduction

With the recent observed Higgs boson candidate at the LHC, the weak sector of the standard model may be approaching closure. However the strong interaction component is far from complete, especially hadronic structure where intense searches are under way for unconventional (non $q\bar{q}$ or qqq) states. Indeed QCD allows for other color singlet combinations, such as glueballs, hybrid mesons, tetraquarks, pentaquarks and dibaryons. As early as 1977 $qq\bar{q}\bar{q}$ tetraquark states were proposed [1] and today there are now good candidates, the $\pi_1(1400)$ and $\pi_1(1600)$ [2,3], listed in the PDG meson summary table [4]. These states have unconventional quantum numbers $J^{PC} = 1^{-+}$ which are not possible for a $q\bar{q}$ system and have been reasonably described [5] as $(q\bar{q})_1(q\bar{q})_1$ color singlet-singlet states referred to as molecules. In the heavy quark sector

Preprint submitted to Elsevier *26 February 2013*

no states with unconventional quantum numbers have yet to be confirmed. However, there are hidden charm states, some also listed in the PDG summary, which can not be described as $c\bar{c}$ mesons. These X, Y and Z particles are charmonium-like but are not consistent with any $c\bar{c}$ spectrum predictions. The lightest is $X(3872)$ which was discovered by Belle in 2003 [6] and has a narrow peak near 3872 MeV in the $\pi^+\pi^- J/\psi$ invariant mass distribution from $B^- \to K^-\pi^+\pi^- J/\psi$ decay. This discovery was confirmed by BaBar in the same decay process [7]. Subsequently, additional X, Y and Z resonances were found in variest experiments and those with confirmed J^{PC} are listed in Table 1.

Table 1
Properties of X, Y and Z mesons with established J^{PC}.

State	Mass (MeV)	$\Gamma(MeV)$	J^{PC}	Decay	Production $B \to$
$X(3872)$	3871.4 ± 0.6	< 2.3	1^{++} or 2^{-+}	$\pi^+\pi^- J/\psi$ $\gamma J/\psi$	$KX(3872)$ $p\bar{p}$
$Z(3930)$	3929 ± 5	29 ± 10	2^{++}	$D\bar{D}$	$\gamma\gamma$
$Y(4008)$	4008^{+82}_{-49}	226^{+97}_{-80}	1^{--}	$\pi^+\pi^- J/\psi$	e^+e^-
$Y(4260)$	4260 ± 12	83 ± 22	1^{--}	$\pi^+\pi^- J/\psi$	e^+e^-
$Y(4360)$	4361 ± 13	74 ± 18	1^{--}	$\pi^+\pi^- \psi'$	e^+e^-
$Y(4660)$	4664 ± 12	48 ± 15	1^{--}	$\pi^+\pi^- \psi'$	e^+e^-

Theoretically, there have been many X, Y and Z investigations using a variety of methods, such as perturbative NRQCD [8,9], lattice QCD [10,11], effective field theory [12,13,14], QCD sum rule [15,16] and potential models [17,18,19,20], all reviewed in detail elsewhere [21,22]. One plausible conjecture [23,24,25] is these states are tetraquarks containing light, $q = u$ or d, and charm, c, quarks in different color schemes. In this paper we investigate this further and consider two color combinations, the somewhat conventional molecular $(q\bar{c})_1(c\bar{q})_1$ singlet-singlet [22] and the more exotic $(qc)_3(\bar{q}\bar{c})_3$ triplet-triplet (diquark) [26]. We reserve the term exotic for the latter since it entails quarks in intermediate color states that are not singlets.

Our work is an extension of a previous light tetraquark study [5] utilizing the QCD Coulomb gauge [CG] model, first implemented to predict a glueball spectrum [27,28] that was in good agreement with lattice QCD data. In this approach the exact QCD Hamiltonian in the Coulomb gauge is approximated by an effective field theoretical relativistic Hamiltonian. The bare parton (current quark and gluon) field operators are dressed by a Bardeen, Cooper and Schrieffer [BCS] rotation and variational ground state (vacuum) minimization. This generates a non-trivial vacuum in which chiral symmetry is dynamically broken and quark/gluon constituent masses and condensates emerge.

The hadrons are represented as quasiparticle excitations using many-body techniques such as Tamm-Dancoff and random phase approximations. This method was subsequently applied to mesons [29,30,31], hybrids [32,33,34,35] and light tetraquark states [5,36].

This paper is organized into five sections. Section 2 specifies the CG model which is then applied to hidden charm tetraquark states in section 3. Numerical results are presented and discussed in section 4 followed by a summary, section 5, detailing conclusions and future work.

2 The QCD Coulomb gauge model

The CG model provides a comprehensive, systematic approach to hadron structure. It is applicable to both quarks and gluons, light and heavy meson and baryon ground and excited states for any flavor and exotic systems involving different combinations of quarks and gluons in various color schemes. It also permits consistent Hamiltonian dynamical mixing between states of entirely different dressed partons while providing insight characteristic of a wave function picture. As further discussed below there are additional attractive theoretical features and it is significant to note that there are no free model parameters as the two dynamical constants, the string tension σ and Coulomb interaction α_s, are predetermined from the literature.

The exact QCD Hamiltonian in the Coulomb gauge [37] is

$$\begin{aligned} H_{\rm QCD} &= H_q + H_g + H_{qg} + H_C & (1)\\ H_q &= \int d\mathbf{x} \Psi^\dagger(\mathbf{x})[-i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla} + \beta m]\Psi(\mathbf{x}) & (2)\\ H_g &= \frac{1}{2}\int d\mathbf{x}\Big[\mathcal{J}^{-1}\boldsymbol{\Pi}^a(\mathbf{x})\cdot\mathcal{J}\boldsymbol{\Pi}^a(\mathbf{x}) + \mathbf{B}^a(\mathbf{x})\cdot\mathbf{B}^a(\mathbf{x})\Big] & (3)\\ H_{qg} &= g\int d\mathbf{x}\, \mathbf{J}^a(\mathbf{x})\cdot\mathbf{A}^a(\mathbf{x}) & (4)\\ H_C &= -\frac{g^2}{2}\int d\mathbf{x}d\mathbf{y}\rho^a(\mathbf{x})\mathcal{J}^{-1}K^{ab}(\mathbf{x},\mathbf{y})\mathcal{J}\rho^b(\mathbf{y})\ , & (5)\end{aligned}$$

where g is the QCD coupling constant, Ψ is the quark field with current quark mass m, $A^a = (\mathbf{A}^a, A_0^a)$ are the gluon fields satisfying the Coulomb (transverse) gauge condition, $\boldsymbol{\nabla}\cdot\mathbf{A^a} = \mathbf{0}$, $(a = 1, 2, ...8)$, $\boldsymbol{\Pi}^a = -\mathbf{E}^a_{tr}$ are the conjugate momenta and

$$\mathbf{E}^a_{tr} = -\dot{\mathbf{A}}^a + g(1 - \nabla^{-2}\boldsymbol{\nabla}\boldsymbol{\nabla}\cdot)f^{abc}A_0^b\mathbf{A}^c \tag{6}$$

$$\mathbf{B}^a = \boldsymbol{\nabla}\times\mathbf{A}^a + \frac{1}{2}gf^{abc}\mathbf{A}^b\times\mathbf{A}^c, \tag{7}$$

are the non-abelian chromodynamic fields. The color densities $\rho^a(\mathbf{x})$ and quark currents $\mathbf{J}^a(\mathbf{x})$ are

$$\rho^a(\mathbf{x}) = \Psi^\dagger(\mathbf{x}) T^a \Psi(\mathbf{x}) + f^{abc}\mathbf{A}^b(\mathbf{x}) \cdot \mathbf{\Pi}^c(\mathbf{x}) \tag{8}$$

$$\mathbf{J}^a(\mathbf{x}) = \Psi^\dagger(\mathbf{x}) \boldsymbol{\alpha} T^a \Psi(\mathbf{x}), \tag{9}$$

where $T^a = \frac{\lambda^a}{2}$ and f^{abc} are the $SU(3)$ color matrices and structure constants, respectively. The Faddeev-Popov determinant, $\mathcal{J} = \det(\mathcal{M})$, of the matrix $\mathcal{M} = \boldsymbol{\nabla} \cdot \mathbf{D}$, with covariant derivative $\mathbf{D}^{ab} = \delta^{ab}\boldsymbol{\nabla} - gf^{abc}\mathbf{A}^c$, is a measure of the gauge manifold curvature and the kernel in Eq. (5) is given by $K^{ab}(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, a|\mathcal{M}^{-1}\nabla^2\mathcal{M}^{-1}|\mathbf{y}, b\rangle$. The Coulomb gauge Hamiltonian is renormalizable, permits resolution of the Gribov problem, preserves rotational invariance, avoids spurious retardation corrections, aids identification of dominant, low energy potentials and does not introduce unphysical degrees of freedom (ghosts).

The bare parton fields have the following normal mode expansions (bare quark spinors u, v, helicity, $\lambda = \pm 1$, and color vectors $\hat{\boldsymbol{\epsilon}}_{\mathcal{C}=1,2,3}$)

$$\Psi(\mathbf{x}) = \int \frac{d\mathbf{k}}{(2\pi)^3} [u_\lambda(\mathbf{k}) b_{\lambda\mathcal{C}}(\mathbf{k}) + v_\lambda(-\mathbf{k}) d^\dagger_{\lambda\mathcal{C}}(-\mathbf{k})] e^{i\mathbf{k}\cdot\mathbf{x}} \hat{\boldsymbol{\epsilon}}_{\mathcal{C}} \tag{10}$$

$$\mathbf{A}^a(\boldsymbol{x}) = \int \frac{d\boldsymbol{k}}{(2\pi)^3} \frac{1}{\sqrt{2k}} [\mathbf{a}^a(\boldsymbol{k}) + \mathbf{a}^{a\dagger}(-\boldsymbol{k})] e^{i\boldsymbol{k}\cdot\boldsymbol{x}} \tag{11}$$

$$\mathbf{\Pi}^a(\boldsymbol{x}) = -i \int \frac{d\boldsymbol{k}}{(2\pi)^3} \sqrt{\frac{k}{2}} [\mathbf{a}^a(\boldsymbol{k}) - \mathbf{a}^{a\dagger}(-\boldsymbol{k})] e^{i\boldsymbol{k}\cdot\boldsymbol{x}}. \tag{12}$$

Our model's starting point is the Coulomb gauge QCD Hamiltonian. In this gauge, the color form of Gauss's law, which is essential for confinement, is satisfied exactly. The unphysical longitudinal degrees of freedom of gluon fields can be completely eliminated by solving Gauss's law. We then make the following approximations: 1) replace the exact Coulomb kernel with a calculable confining potential; 2) use the lowest order, unit value for the Faddeev-Popov determinant. Therefore, H_C in the CG model Hamiltonian becomes

$$H_C^{\mathrm{CG}} = -\frac{1}{2} \int d\mathbf{x} d\mathbf{y} \rho^a(\mathbf{x}) \hat{V}(|\mathbf{x} - \mathbf{y}|) \rho^a(\mathbf{y}). \tag{13}$$

The confining and leading canonical interaction is represented by a Cornell type potential $\hat{V}(r) = -\alpha_s/r + \sigma r$. Previous studies with this interaction were in good agreement with both lattice glueball masses [27] and the observed meson spectrum [30]. Performing a fourier transform, the potential in momentum space is

$$V(|\mathbf{k}|) = -\frac{4\pi\alpha_s}{k^2} - \frac{8\pi\sigma}{k^4}. \tag{14}$$

Next, hadron states are expressed as BCS vacuum $|\Omega\rangle$ excitations involving dressed quark (antiquark) Fock operators $B^{\dagger}_{\lambda\mathcal{C}}$, $D_{\lambda\mathcal{C}}$ related to the bare operators $b_{\lambda\mathcal{C}}$, $d^{\dagger}_{\lambda\mathcal{C}}$ by the rotation

$$B_{\lambda\mathcal{C}}(\mathbf{k}) = \cos\frac{\theta(k)}{2} b_{\lambda\mathcal{C}}(\mathbf{k}) - \lambda\sin\frac{\theta(k)}{2} d^{\dagger}_{\lambda\mathcal{C}}(-\mathbf{k}) \tag{15}$$

$$D_{\lambda\mathcal{C}}(-\mathbf{k}) = \cos\frac{\theta(k)}{2} d_{\lambda\mathcal{C}}(-\mathbf{k}) + \lambda\sin\frac{\theta(k)}{2} b^{\dagger}_{\lambda\mathcal{C}}(\mathbf{k}), \tag{16}$$

$$\tag{17}$$

where $\theta(k)$ is the BCS angle. Correspondingly, the dressed Dirac spinors are

$$\mathcal{U}_{\lambda}(\mathbf{k}) = \frac{1}{\sqrt{2}}\begin{pmatrix} \sqrt{1+\sin\phi(k)}\chi_{\lambda} \\ \sqrt{1-\sin\phi(k)}\boldsymbol{\sigma}\cdot\hat{\mathbf{k}}\chi_{\lambda} \end{pmatrix} \tag{18}$$

$$\mathcal{V}_{\lambda}(-\mathbf{k}) = \frac{1}{\sqrt{2}}\begin{pmatrix} -\sqrt{1-\sin\phi(k)}\boldsymbol{\sigma}\cdot\hat{\mathbf{k}}\chi_{\lambda} \\ \sqrt{1+\sin\phi(k)}\chi_{\lambda} \end{pmatrix}, \tag{19}$$

where $\phi(k)$ is the gap angle from the gap equation that minimizes the energy of the BCS vacuum [34] and is related to $\theta(k)$ by $tan(\phi(k)-\theta(k)) = m/k$.

The four model parameters have all been predetermined [31]. The light and charm bare quark masses are 5 MeV and 1350 MeV, respectively, while the two dynamic constants are $\sigma = 0.18\ GeV^2$ and $\alpha_s = \frac{g^2}{4\pi} = 0.4$. This few parameter, constrained model has successfully described meson, hybrid, glueball and light tetraquark systems yielding results agreeing with experimental and lattice data.

3 Application to hidden charm tetraquark states

For this heavier tetraquark system, the quark (antiquark) cm momenta are $\mathbf{k}_{i=1,2,3,4}$ and the following wave function ansatz is adopted

$$|\Psi^{JPC}\rangle = \int d\mathbf{k}_1 d\mathbf{k}_2 d\mathbf{k}_3 d\mathbf{k}_4 \delta(\mathbf{k}_1+\mathbf{k}_2+\mathbf{k}_3+\mathbf{k}_4)\Phi^{JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_i) \times R^{\mathcal{C}_1\mathcal{C}_2}_{\mathcal{C}_3\mathcal{C}_4} B^{\dagger}_{\lambda_1\mathcal{C}_1}(\mathbf{k}_1) B^{\dagger}_{\lambda_2\mathcal{C}_2}(\mathbf{k}_2) D^{\dagger}_{\lambda_3\mathcal{C}_3}(\mathbf{k}_3) D^{\dagger}_{\lambda_4\mathcal{C}_4}(\mathbf{k}_4)|\Omega\rangle. \tag{20}$$

The expression for the matrix $R^{\mathcal{C}_1\mathcal{C}_2}_{\mathcal{C}_3\mathcal{C}_4}$ depends on the specific color scheme selected. As depicted in Fig. 1, the $SU_c(3)$ color algebra for four quarks produces 81 color states, $3\otimes\bar{3}\otimes3\otimes\bar{3} = 27\oplus10\oplus\bar{10}\oplus8\oplus8\oplus8\oplus8\oplus1\oplus1$ of which two



Fig. 1. Four tetraquark color schemes. One is a singlet-singlet molecule while the other three are more exotic atoms (octet and two diquark schemes).

are overall color singlets that can be obtained in four different schemes (only two are independent) depending on the intermediate coupling of two quarks: singlet-singlet (molecule), octet-octet and two diquark schemes triplet-triplet and sextet-sextet. In this article we focus on singlet-singlet (molecule) and triplet-triplet (diquark) schemes, which provide the lightest states.

For the diquark configuration, the wave function is

$$\Phi^{JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_i) = F^{JPC}(\mathbf{k}_i)\langle\frac{1}{2}\frac{1}{2}\lambda_1\lambda_2|S_A\lambda_A\rangle\langle\frac{1}{2}\frac{1}{2}\lambda_3\lambda_4|S_B\lambda_B\rangle\langle S_AS_B\lambda_A\lambda_B|S\lambda\rangle$$
$$\langle L_AL_Bm_Am_B|lm\rangle\langle lL_Imm_I|LM\rangle\langle SL\lambda M|JM_J\rangle Y^{m_A}_{L_A}(\hat{\mathbf{k}}_A)Y^{m_B}_{L_B}(\hat{\mathbf{k}}_B)Y^{m_I}_{L_I}(\hat{\mathbf{k}}_I),$$

where $Y^m_L(\hat{\mathbf{k}})$ is the spherical harmonic function, while $S_{A(B)}$, $L_{A(B)}$ and L_I are the $A(B)$ diquark total spin, orbit angular momentum and orbit angular momentum between the two diquarks, respectively. The radial wave function, $F^{JPC}(\mathbf{k}_i)$, is chosen to be a Gaussian, $\exp(-\frac{k_A^2}{\alpha_A^2}-\frac{k_B^2}{\alpha_B^2}-\frac{k_I^2}{\alpha_I^2})$, where $\mathbf{k}_A$, $\mathbf{k}_B$ and $\mathbf{k}_I$ are respectively $\mathbf{k}_A=\frac{\mathbf{k}_1-\mathbf{k}_2}{2}$, $\mathbf{k}_B=\frac{\mathbf{k}_3-\mathbf{k}_4}{2}$, $\mathbf{k}_I=\frac{\mathbf{k}_1+\mathbf{k}_2}{2}-\frac{\mathbf{k}_3+\mathbf{k}_4}{2}$ and α_A, α_B and α_I are the variational parameters.

The variational mass is then given by

$$M_{JPC} = \frac{\langle\Psi^{JPC}|H^{CG}|\Psi^{JPC}\rangle}{\langle\Psi^{JPC}|\Psi^{JPC}\rangle} = M_{self} + M_{qq} + M_{q\bar{q}} + M_{\bar{q}\bar{q}} + M_{annih}\ ,$$

where the subscripts indicate the contribution from the single particle quark self energy and two particle qq, $q\bar{q}$, $\bar{q}\bar{q}$ scattering and $q\bar{q}$ annihilation, respectively. All contributions to the tetraquark mass are diagramed in Fig. 2.

Using the above specified Hamiltonian and wave function, the self energy term can be reduced to



Fig. 2. Equal-time diagrams for tetraquarks. The first four diagrams are for the kinetic and self energy. The last diagram is for the annihilation term and the others are for the qq, $q\bar{q}$ and $\bar{q}\bar{q}$ interactions.

$$M_{self} = \sum_{i=1}^{4} \int \frac{d\mathbf{k}_1}{(2\pi)^3} \frac{d\mathbf{k}_2}{(2\pi)^3} \frac{d\mathbf{k}_3}{(2\pi)^3} \frac{d\mathbf{k}_4}{(2\pi)^3} \Phi^{\dagger JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3,\mathbf{k}_4) \\ \times E_i(\mathbf{k}_i)\Phi^{JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3,\mathbf{k}_4), \tag{21}$$

where the single particle energy is given by

$$E_i(\mathbf{k}_i) = m_i sin\,\theta(\mathbf{k}_i) + k_i cos\,\theta(\mathbf{k}_i) - \frac{2}{3}\int \frac{d\mathbf{q}}{(2\pi)^3} V(|\mathbf{q}|)[sin\,\theta(\mathbf{k}_i)\sin\,\theta(\mathbf{q}) \\ + cos\,\theta(\mathbf{k}_i) cos\,\theta(\mathbf{q})\hat{\mathbf{q}}\cdot\hat{\mathbf{k}}_i]. \tag{22}$$

Similarly, the two particle contributions can be evaluated yielding

$$M_{qq} = -\frac{2}{3}\int \frac{d\mathbf{k}_A}{(2\pi)^3} \frac{d\mathbf{k}_B}{(2\pi)^3} \frac{d\mathbf{k}_I}{(2\pi)^3} \frac{d\mathbf{q}}{(2\pi)^3} \mathcal{U}^{\dagger}_{\lambda_1'}(\mathbf{k}_1')\mathcal{U}_{\lambda_1}(\mathbf{k}_1)\mathcal{U}^{\dagger}_{\lambda_2'}(\mathbf{k}_2')\mathcal{U}_{\lambda_2}(\mathbf{k}_2) \\ \times V(|\mathbf{q}|)\Phi^{\dagger JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3,\mathbf{k}_4)\Phi^{JPC}_{\lambda_1'\lambda_2'\lambda_3'\lambda_4'}(\mathbf{k}_1',\mathbf{k}_2',\mathbf{k}_3',\mathbf{k}_4'), \tag{23}$$

$$M_{q\bar{q}} = \frac{1}{3}\int \frac{d\mathbf{k}_A}{(2\pi)^3} \frac{d\mathbf{k}_B}{(2\pi)^3} \frac{d\mathbf{k}_I}{(2\pi)^3} \frac{d\mathbf{q}}{(2\pi)^3} \mathcal{U}^{\dagger}_{\lambda_1'}(\mathbf{k}_1')\mathcal{U}_{\lambda_1}(\mathbf{k}_1)\mathcal{V}^{\dagger}_{\lambda_3}(\mathbf{k}_3)\mathcal{V}_{\lambda_3'}(\mathbf{k}_3') \\ \times V(|\mathbf{q}|)\Phi^{\dagger JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3,\mathbf{k}_4)\Phi^{JPC}_{\lambda_1'\lambda_2'\lambda_3'\lambda_4'}(\mathbf{k}_1',\mathbf{k}_2',\mathbf{k}_3',\mathbf{k}_4'), \tag{24}$$

$$M_{\bar{q}\bar{q}} = -\frac{2}{3}\int \frac{d\mathbf{k}_A}{(2\pi)^3} \frac{d\mathbf{k}_B}{(2\pi)^3} \frac{d\mathbf{k}_I}{(2\pi)^3} \frac{d\mathbf{q}}{(2\pi)^3} \mathcal{V}^{\dagger}_{\lambda_1'}(\mathbf{k}_1')\mathcal{V}_{\lambda_1}(\mathbf{k}_1)\mathcal{V}^{\dagger}_{\lambda_2'}(\mathbf{k}_2')\mathcal{V}_{\lambda_2}(\mathbf{k}_2) \\ \times V(|\mathbf{q}|)\Phi^{\dagger JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3,\mathbf{k}_4)\Phi^{JPC}_{\lambda_1'\lambda_2'\lambda_3'\lambda_4'}(\mathbf{k}_1',\mathbf{k}_2',\mathbf{k}_3',\mathbf{k}_4'), \tag{25}$$

where $\mathbf{q}$ is the momentum transfer $\mathbf{k}_1' - \mathbf{k}_1$. Finally, the annihilation term is

$$M_{annih} = \frac{1}{3}\int \frac{d\mathbf{k}_A}{(2\pi)^3}\frac{d\mathbf{k}_B}{(2\pi)^3}\frac{d\mathbf{k}_I}{(2\pi)^3}\frac{d\mathbf{q}}{(2\pi)^3} \times \mathcal{V}^\dagger_{\lambda_4}(\mathbf{k}_4)\mathcal{U}_{\lambda_1}(\mathbf{k}_1)\mathcal{U}^\dagger_{\lambda'_1}(\mathbf{k}'_1)\mathcal{V}_{\lambda'_4}(\mathbf{k}'_4)$$
$$\times V(|\mathbf{q}||)\Phi^{\dagger JPC}_{\lambda_1\lambda_2\lambda_3\lambda_4}(\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3,\mathbf{k}_4)\Phi^{JPC}_{\lambda'_1\lambda'_2\lambda'_3\lambda'_4}(\mathbf{k}'_1,\mathbf{k}'_2,\mathbf{k}'_3,\mathbf{k}'_4), \qquad (26)$$

where now $\mathbf{q}$ is the annihilation momentum $\mathbf{k}_1 + \mathbf{k}_4$. For the molecular case, each $q\bar{q}$ pair forms a subgroup and the corresponding wave function and mass formula are similar. The 12-dimension integrals are performed numerically using the Monte-Carlo method.

4 Numerical Results

We calculated all possible states with $S_i = 0, 1$ and/or $L_i = 0, 1$ and the numerical results for corresponding J^{PC} are listed in Table 2. For model completeness and further comparative insight, we also list our $c\bar{c}g$ hybrid predictions from an earlier publication [34]. It is clear that the diquark composition is heavier than the corresponding molecular state. Even more significant is the sensitivity to arrangement of flavor as $(q\bar{q})_1(c\bar{c}_1)$ states are much lighter than $(q\bar{c})_1(\bar{q}c)_1$ for the same L_i, S_i configuration. The CG model therefore predicts that the lightest tetraquark states are molecules having quarks with the same flavor in a color singlet (i.e. a $\eta\eta_c$ or $\omega J/\psi$ type flavor structure depending upon spin).

For the s wave triplet-triplet tetraquark states, the mass changes little with different spin. This is because there is a heavy charm quark in each color triplet

Table 2
Predicted spectrum for tetraquarks (singlet-singlet and triplet-triplet) and hybrid mesons in MeV. Unless explicitly stated, all L_i and S_i are zero.

L_i, S_i	J^{PC}	$(q\bar{q})_1(c\bar{c})_1$	J^{PC}	$(q\bar{c})_1(c\bar{q})_1$	$(qc)_{\bar{3}}(\bar{c}\bar{q})_3$	J^{PC}	$(c\bar{c})_8 g$
$L_i, S_i = 0$	0^{++}	3540	0^{++}	4129	4428	0^{++}	3945
$L_I = 1$	1^{-+}	3540	1^{--}	4129	4741	1^{-+}	4020
$L_A = 1$	1^{--}	4186	1^{-+}	4563	4826	1^{-+}	4155
$L_B = 1$	1^{--}	4026	1^{-+}	4565	4825	1^{-+}	4565
$S_A = 1$	1^{+-}	3781	1^{++}	4137	4430	1^{++}	4100
$S_B = 1$	1^{+-}	3538	1^{++}	4137	4430	1^{+-}	3830
$S_A, S_B = 1$	0^{++}	3783	0^{++}	4144	4405	0^{--}	4020
$S_A, S_B = 1$	1^{++}	3783	1^{++}	4144	4419	3^{-+}	4615
$S_A, S_B = 1$	2^{++}	3783	2^{++}	4144	4444	2^{++}	3965

and the spin splitting decreases with heavier quark mass. For orbit angular momentum 1, the triplet-triplet tetraquark mass increases about 300 to 400 MeV and all p wave tetraquarks have mass above 4400 MeV.

Similarly for the $(q\bar{c})_1(c\bar{q})_1$ states, the spin splitting is very small due to the large charm quark mass in each color singlet, and again if either L_A or L_B changes from 0 to 1 the mass increases, now about 430 MeV. For the $(q\bar{q})_1(c\bar{c})_1$ flavor arrangement, the mass does not depend on the $c\bar{c}$ spin but increases about 240 MeV if the spin of light quark subgroup changes from 0 to 1. The mass increase due to larger orbit angular momentum is also flavor sensitive, since for $q\bar{q}$ p waves it is about 650 MeV, while for $c\bar{c}$ p waves it is 490 MeV.

In Fig. 3, we compare predicted 1^{++} and 1^{--} tetraquark masses to observed X and Y states. Only about half of the states have established J^{PC} and to aid determining quantum numbers it is noteworthy that the observed $Y(4140)$, $X(4160)$, $Y(4274)$ and $X(4350)$ are comparable to the heaviest model 1^{++} states. If this identification proves correct, then the model predicts the $Y(4140)$ and $X(4160)$ are molecular states consisting of two spin 0 clusters having a $D\bar{D}$ flavor composition while the $Y(4274)$ and $X(4350)$ would be exotic spin 1 triplet-triplet states with a diquark composition.

Regarding the established J^{PC} states, the predicted $(q\bar{q})_1(c\bar{c})_1$ 1^{++} mass is 3783 MeV about 3% less than observed, suggesting the $X(3872)$ may be a molecular state with a $\omega J/\psi$ flavor composition. Four 1^{--} tetraquark states are predicted with the lightest mass at 4026 MeV which is close to $Y(4008)$ discovered by Belle [38]. The masses of the next two heavier molecular states



Fig. 3. Theoretical results (dashed lines) for 1^{++} and 1^{--} tetraquark states compared to data (solid lines).

are 4129 and 4186, which are comparable with $Y(4260)$ and $Y(4360)$. The states with masses 4026 MeV and 4186 MeV have a $\eta\eta_c$ flavor arrangement while the state at 4129 MeV is a $D\bar{D}$ type molecule. The calculated mass of the triplet-triplet 1^{--} state is 4741 MeV which is also close to the $Y(4660)$ mass. This state would be an exotic diquark if this assignment prevails. Lastly, for the observed 2^{++} $Z(3930)$ listed in Table 1, the model tetraquark states are either above or below this mass by over 150 MeV, however the predicted $c\bar{c}g$ hybrid mass of 3965 MeV (see Table 2) agrees quite well. Similarly the 0^{++} hybrid at 3945 MeV is very close to the $X(3915)$. This state does not have established quantum numbers other than the $G = +1$ parity which is also the same as our hybrid prediction. The hybrid results are quite firm since they include both hyperfine and the non-linear (non-Abelian) component of the color magnetic fields (Eq. (7)).

Finally, it is interesting that the decay channels summarized in Table 1 are consistent with the Coulomb gauge model flavor structure predictions for those observed states. Specifically, for the predicted molecular state assignments involving the X(3872), Y(4008) and Y(4260) having a $\omega J/\psi$ flavor structure, their decays would favor the observed $\pi^{\pm} J/\psi$ channel. For the diquark model states, detailed calculations would be necessary to make firm predictions. Similarly, for the undetermined J^{PC} states predicted to be $D\bar{D}$ molecules, $Y(4140)$ and $X(4160)$, their decays would would favor the $D\bar{D}$ channel and, if observed, would be an identifying signature.

5 Summary

We have applied the Coulomb gauge model to tetraquark systems with hidden charm having two different color compositions, singlet-singlet and triplet-triplet. Generally, triplet-triplet states are the heaviest for the same spin or orbit angular momentum. For color singlet-singlet states, we further investigated two flavor groupings corresponding to $\eta\eta_c$ or J/ψ versus $D\bar{D}$ or $D^*\bar{D}^*$ type molecules (meson-meson states) and found that the same flavor color singlets produce a lower mass. Also, the mass from the spin part decreases rapidly with increasing quark mass, while the orbit angular momentum contribution decreases slowly with increasing quark mass.

Our key finding is the $X(3872)$ mass is close to the predicted 1^{++} $(q\bar{q})_1(c\bar{c})_1$ state suggesting it is a conventional $\omega J/\psi$ type molecule and not a more exotic diquark tetraquark state. For 1^{--}, four states are predicted, three molecular and one triplet-triplet. Their masses are comparable with $Y(4008)$, $Y(4260)$, $Y(4360)$ and $Y(4660)$. The masses of other states with undetermined J^{PC} are close to the calculated 1^{++} results. Therefore, the J^{PC} number of these states could be 1^{++}.

Future work will address several issues. The spin dependence will be further studied now including the hyperfine interaction [31]. Also, a dynamic mixing analysis will be performed involving $q\bar{q}$ and $qq\bar{q}\bar{q}$ states having various flavor distributions as well as mixing with glueball and hybrid meson exotic states. Finally, the CG model will be applied to b quark systems to aid experimental searched at higher energies.

Acknowledgement

The authors are deeply grateful to Felipe Llanes-Estrada for insightful comments. This work is supported in part by DFG and NSFC (CRC 110) and by National Natural Science Foundation of China (Grant No. 11035006).

References

[1] R. L. Jaffe, Phys. Rev. D **15** (1977) 267.

[2] G. S. Adams et al. [E852 Collaboration], Phys. Rev. Lett. **81** (1998) 5760.

[3] S. U. Chung et al. [E852 Collaboration], Phys. Rev. D **60** (1999) 092001 [hep-ex/9902003].

[4] J. Beringer et al. (Particle Data Group), Phys. Rev. D **86** (2012) 010001.

[5] I. J. General, P. Wang, S. R. Cotanch and F. J. Llanes-Estrada, Phys. Lett. B **653** (2007) 216 [arXiv:0707.1286 [hep-ph]].

[6] S. K. Choi et al. [Belle Collaboration], Phys. Rev. Lett. **91** (2003) 262001 [hep-ex/0309032].

[7] B. Aubert et al. [BaBar Collaboration], Phys. Rev. D **71** (2005) 031501 [hep-ex/0412051].

[8] G. T. Bodwin, E. Braaten and G. P. Lepage, Phys. Rev. D **51** (1995) 1125 [Erratum-ibid. D **55** (1997) 5853] [hep-ph/9407339].

[9] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Nucl. Phys. B **566** (2000) 275 [hep-ph/9907240].

[10] M. Okamoto et al. [CP-PACS Collaboration], Phys. Rev. D **65** (2002) 094508 [hep-lat/0112020].

[11] J. J. Dudek, R. G. Edwards, N. Mathur and D. G. Richards, Phys. Rev. D **77** (2008) 034501 [arXiv:0707.4162 [hep-lat]].

[12] E. Braaten and M. Lu, Phys. Rev. D **74** (2006) 054020 [hep-ph/0606115].

[13] E. Braaten and M. Kusunoki, Phys. Rev. D **72** (2005) 014012 [hep-ph/0506087].

[14] D. L. Canham, H. -W. Hammer and R. P. Springer, Phys. Rev. D **80** (2009) 014009 [arXiv:0906.1263 [hep-ph]].

[15] R. D'E. Matheus, S. Narison, M. Nielsen and J. M. Richard, Phys. Rev. D **75** (2007) 014005 [hep-ph/0608297].

[16] S. H. Lee, M. Nielsen and U. Wiedner, arXiv:0803.1168 [hep-ph].

[17] X. Liu, Z. -G. Luo, Y. -R. Liu and S. -L. Zhu, Eur. Phys. J. C **61** (2009) 411 [arXiv:0808.0073 [hep-ph]].

[18] L. Cao, Y. -C. Yang and H. Chen, Few Body Syst. **53** (2012) 327 [arXiv:1206.3008 [hep-ph]].

[19] B. -Q. Li and K. -T. Chao, Phys. Rev. D **79** (2009) 094004 [arXiv:0903.5506 [hep-ph]].

[20] Bhaghyesh and K. B. Vijaya Kumar, Commun. Theor. Phys. **55** (2011) 1044.

[21] S. Godfrey and S. L. Olsen, Ann. Rev. Nucl. Part. Sci. **58** (2008) 51 [arXiv:0801.3867 [hep-ph]].

[22] E. S. Swanson, Phys. Rept. **429** (2006) 243 [hep-ph/0601110].

[23] E. S. Swanson, Phys. Lett. B **588** (2004) 189 [hep-ph/0311229].

[24] F. E. Close and P. R. Page, Phys. Lett. B **578** (2004) 119 [hep-ph/0309253].

[25] M. B. Voloshin, Phys. Lett. B **579** (2004) 316 [hep-ph/0309307].

[26] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. D **71** (2005) 014028 [hep-ph/0412098].

[27] A. P. Szczepaniak, E. S. Swanson, C. -R. Ji and S. R. Cotanch, Phys. Rev. Lett. **76** (1996) 2011 [hep-ph/9511422].

[28] S. R. Cotanch, A. P. Szczepaniak, E. S. Swanson and C. R. Ji, Nucl. Phys. A **631** (1998) 640C.

[29] F. J. Llanes-Estrada and S. R. Cotanch, Phys. Rev. Lett. **84** 1102 (2000) [hep-ph/9906359].

[30] F. J. Llanes-Estrada and S. R. Cotanch, Nucl. Phys. A **697** (2002) 303 [hep-ph/0101078].

[31] F. J. Llanes-Estrada, S. R. Cotanch, A. P. Szczepaniak and E. S. Swanson, Phys. Rev. C **70** (2004) 035202.

[32] S. R. Cotanch and F. J. Llanes-Estrada, Nucl. Phys. A **689** (2001) 481.

[33] F. J. Llanes-Estrada and S. R. Cotanch, Phys. Lett. B **504** (2001) 15 [hep-ph/0008337].

[34] I. J. General, S. R. Cotanch and F. J. Llanes-Estrada, Eur. Phys. J. C **51** (2007) 347 [hep-ph/0609115].

[35] S. R. Cotanch, I. J. General and P. Wang, Eur. Phys. J. A **31** (2007) 656.

[36] P. Wang, S. R. Cotanch and I. J. General, Eur. Phys. J. C **55** (2008) 409 [arXiv:0801.4810 [hep-ph]].

[37] T. D. Lee, Particle Physics and Introduction to Field Theory, Harwood Academic Publishers, New York (1990).

[38] C. Z. Yuan et al. [Belle Collaboration], Phys. Rev. Lett. **99** (2007) 182004 [arXiv:0707.2541 [hep-ex]].